

| | | | |
|---------------------|--|------------------|--|
| Surname | | Other Names | |
| Centre Number | | Candidate Number | |
| Candidate Signature | | | |

| |
|--------------------|
| For Examiner's Use |
|--------------------|

General Certificate of Education
June 2007
Advanced Level Examination



PHYSICS (SPECIFICATION A)
Unit 6 Nuclear Instability: Medical Physics Option

PHA6/W

Thursday 14 June 2007 9.00 am to 10.15 am

| |
|---|
| <p>For this paper you must have:</p> <ul style="list-style-type: none"> • a calculator • a pencil and a ruler. |
|---|

Time allowed: 1 hour 15 minutes

Instructions

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Answer the questions in the spaces provided.
- Show all your working.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The maximum mark for this paper is 40.
- Two of these marks will be awarded for using good English, organising information clearly and using specialist vocabulary where appropriate.
- The marks for questions are shown in brackets.
- A *Data sheet* is provided on pages 3 and 4. You may wish to detach this perforated sheet at the start of the examination.
- You are expected to use a calculator where appropriate.
- Questions 1(a) and 5(a) should be answered in continuous prose. In these questions you will be marked on your ability to use good English, to organise information clearly and to use specialist vocabulary where appropriate.

| For Examiner's Use | | | |
|----------------------------------|------|----------|------|
| Question | Mark | Question | Mark |
| 1 | | | |
| 2 | | | |
| 3 | | | |
| 4 | | | |
| 5 | | | |
| | | | |
| | | | |
| | | | |
| Total (Column 1) | | → | |
| Total (Column 2) | | → | |
| Quality of Written Communication | | | |
| TOTAL | | | |
| Examiner's Initials | | | |

Data Sheet

- A perforated *Data Sheet* is provided as pages 3 and 4 of this question paper.
- This sheet may be useful for answering some of the questions in the examination.
- You may wish to detach this sheet before you begin work.

Data Sheet

| Fundamental constants and values | | | | Mechanics and Applied Physics | | Fields, Waves, Quantum Phenomena | |
|--|----------------|-------------------------|-----------------------------------|--|---|----------------------------------|--|
| Quantity | Symbol | Value | Units | | | | |
| speed of light in vacuo | c | 3.00×10^8 | m s^{-1} | $v = u + at$ | $g = \frac{F}{m}$ | | |
| permeability of free space | μ_0 | $4\pi \times 10^{-7}$ | H m^{-1} | $s = \left(\frac{u+v}{2}\right)t$ | $g = -\frac{GM}{r^2}$ | | |
| permittivity of free space | ϵ_0 | 8.85×10^{-12} | F m^{-1} | $s = ut + \frac{at^2}{2}$ | $g = -\frac{\Delta V}{\Delta x}$ | | |
| charge of electron | e | 1.60×10^{-19} | C | $v^2 = u^2 + 2as$ | $V = -\frac{GM}{r}$ | | |
| the Planck constant | h | 6.63×10^{-34} | J s | $F = \frac{\Delta(mv)}{\Delta t}$ | $a = -(2\pi f)^2 x$ | | |
| gravitational constant | G | 6.67×10^{-11} | $\text{N m}^2 \text{kg}^{-2}$ | $P = Fv$ | $v = \pm 2\pi f \sqrt{A^2 - x^2}$ | | |
| the Avogadro constant | N_A | 6.02×10^{23} | mol^{-1} | $\text{efficiency} = \frac{\text{power output}}{\text{power input}}$ | $x = A \cos 2\pi ft$ | | |
| molar gas constant | R | 8.31 | $\text{J K}^{-1} \text{mol}^{-1}$ | $\omega = \frac{v}{r} = 2\pi f$ | $T = 2\pi \sqrt{\frac{m}{k}}$ | | |
| the Boltzmann constant | k | 1.38×10^{-23} | J K^{-1} | $a = \frac{v^2}{r} = r\omega^2$ | $T = 2\pi \sqrt{\frac{l}{g}}$ | | |
| the Stefan constant | σ | 5.67×10^{-8} | $\text{W m}^{-2} \text{K}^{-4}$ | $I = \sum mr^2$ | $\lambda = \frac{\omega s}{D}$ | | |
| the Wien constant | a | 2.90×10^{-3} | m K | $E_k = \frac{1}{2} I\omega^2$ | $d \sin \theta = n\lambda$ | | |
| electron rest mass | m_e | 9.11×10^{-31} | kg | $\omega_2 = \omega_1 + \alpha t$ | $\theta \approx \frac{\lambda}{D}$ | | |
| (equivalent to $5.5 \times 10^{-4} \text{u}$) | | | | $\theta = \omega_1 t + \frac{1}{2} \alpha t^2$ | ${}^1n_2 = \frac{\sin \theta_1}{\sin \theta_2} = \frac{c_1}{c_2}$ | | |
| electron charge/mass ratio | e/m_e | 1.76×10^{11} | C kg^{-1} | $\omega_2^2 = \omega_1^2 + 2\alpha\theta$ | ${}^1n_2 = \frac{n_2}{n_1}$ | | |
| proton rest mass | m_p | 1.67×10^{-27} | kg | $\theta = \frac{1}{2} (\omega_1 + \omega_2)t$ | $\sin \theta_c = \frac{1}{n}$ | | |
| (equivalent to 1.00728u) | | | | $T = I\alpha$ | $E = hf$ | | |
| proton charge/mass ratio | e/m_p | 9.58×10^7 | C kg^{-1} | $\text{angular momentum} = I\omega$ | $hf = \phi + E_k$ | | |
| neutron rest mass | m_n | 1.67×10^{-27} | kg | $W = T\theta$ | $hf = E_1 - E_2$ | | |
| (equivalent to 1.00867u) | | | | $P = T\omega$ | $\lambda = \frac{h}{p} = \frac{h}{mv}$ | | |
| gravitational field strength | g | 9.81 | N kg^{-1} | $\text{angular impulse} = \text{change of angular momentum} = Tt$ | $c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$ | | |
| acceleration due to gravity | g | 9.81 | m s^{-2} | $\Delta Q = \Delta U + \Delta W$ | | | |
| atomic mass unit | u | 1.661×10^{-27} | kg | $\Delta W = p\Delta V$ | | | |
| (1u is equivalent to 931.3 MeV) | | | | $pV^\gamma = \text{constant}$ | | | |
| Fundamental particles | | | | $\text{work done per cycle} = \text{area of loop}$ | Electricity | | |
| Class | Name | Symbol | Rest energy /MeV | $\text{input power} = \text{calorific value} \times \text{fuel flow rate}$ | $\epsilon = \frac{E}{Q}$ | | |
| photon | photon | γ | 0 | $\text{indicated power as (area of } p-V \text{ loop)} \times (\text{no. of cycles/s}) \times (\text{no. of cylinders})$ | $\epsilon = I(R+r)$ | | |
| lepton | neutrino | ν_e | 0 | $\text{friction power} = \text{indicated power} - \text{brake power}$ | $\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$ | | |
| | | ν_μ | 0 | $\text{efficiency} = \frac{W}{Q_{in}} = \frac{Q_{in} - Q_{out}}{Q_{in}}$ | $R_T = R_1 + R_2 + R_3 + \dots$ | | |
| | electron | e^\pm | 0.510999 | $\text{maximum possible efficiency} = \frac{T_H - T_C}{T_H}$ | $P = I^2 R$ | | |
| | muon | μ^\pm | 105.659 | | $E = \frac{F}{Q} = \frac{V}{d}$ | | |
| mesons | pion | π^\pm | 139.576 | | $E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$ | | |
| | | π^0 | 134.972 | | $E = \frac{1}{2} QV$ | | |
| | kaon | K^\pm | 493.821 | | $F = BIl$ | | |
| | | K^0 | 497.762 | | $F = BQv$ | | |
| baryons | proton | p | 938.257 | | $Q = Q_0 e^{-t/RC}$ | | |
| | neutron | n | 939.551 | | $\Phi = BA$ | | |
| Properties of quarks | | | | | | | |
| Type | Charge | Baryon number | Strangeness | | | | |
| u | $+\frac{2}{3}$ | $+\frac{1}{3}$ | 0 | | | | |
| d | $-\frac{1}{3}$ | $+\frac{1}{3}$ | 0 | | | | |
| s | $-\frac{1}{3}$ | $+\frac{1}{3}$ | -1 | | | | |
| Geometrical equations | | | | | | | |
| arc length = $r\theta$ | | | | | | | |
| circumference of circle = $2\pi r$ | | | | | | | |
| area of circle = πr^2 | | | | | | | |
| area of cylinder = $2\pi rh$ | | | | | | | |
| volume of cylinder = $\pi r^2 h$ | | | | | | | |
| area of sphere = $4\pi r^2$ | | | | | | | |
| volume of sphere = $\frac{4}{3} \pi r^3$ | | | | | | | |

Turn over ►

$$\text{magnitude of induced emf} = N \frac{\Delta\Phi}{\Delta t}$$

$$I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$$

$$V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$$

Mechanical and Thermal Properties

$$\text{the Young modulus} = \frac{\text{tensile stress}}{\text{tensile strain}} = \frac{F}{A} \frac{l}{e}$$

$$\text{energy stored} = \frac{1}{2} Fe$$

$$\Delta Q = mc \Delta\theta$$

$$\Delta Q = ml$$

$$pV = \frac{1}{3} Nmc^2$$

$$\frac{1}{2} mc^2 = \frac{3}{2} kT = \frac{3RT}{2N_A}$$

Nuclear Physics and Turning Points in Physics

$$\text{force} = \frac{eV_p}{d}$$

$$\text{force} = Bev$$

$$\text{radius of curvature} = \frac{mv}{Be}$$

$$\frac{eV}{d} = mg$$

$$\text{work done} = eV$$

$$F = 6\pi\eta rv$$

$$I = k \frac{I_0}{x^2}$$

$$\frac{\Delta N}{\Delta t} = -\lambda N$$

$$\lambda = \frac{h}{\sqrt{2meV}}$$

$$N = N_0 e^{-\lambda t}$$

$$T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$$

$$R = r_0 A^{\frac{1}{3}}$$

$$E = mc^2 = \frac{m_0 c^2}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

$$l = l_0 \left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}$$

$$t = \frac{t_0}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

Astrophysics and Medical Physics

| Body | Mass/kg | Mean radius/m |
|-------|-----------------------|--------------------|
| Sun | 2.00×10^{30} | 7.00×10^8 |
| Earth | 6.00×10^{24} | 6.40×10^6 |

$$1 \text{ astronomical unit} = 1.50 \times 10^{11} \text{ m}$$

$$1 \text{ parsec} = 206265 \text{ AU} = 3.08 \times 10^{16} \text{ m} = 3.26 \text{ ly}$$

$$1 \text{ light year} = 9.45 \times 10^{15} \text{ m}$$

$$\text{Hubble constant } (H) = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$$

$$M = \frac{\text{angle subtended by image at eye}}{\text{angle subtended by object at unaided eye}}$$

$$M = \frac{f_o}{f_e}$$

$$m - M = 5 \log \frac{d}{10}$$

$$\lambda_{\text{max}} T = \text{constant} = 0.0029 \text{ m K}$$

$$v = Hd$$

$$P = \sigma AT^4$$

$$\frac{\Delta f}{f} = \frac{v}{c}$$

$$\frac{\Delta \lambda}{\lambda} = -\frac{v}{c}$$

$$R_s \approx \frac{2GM}{c^2}$$

Medical Physics

$$\text{power} = \frac{1}{f}$$

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f} \text{ and } m = \frac{v}{u}$$

$$\text{intensity level} = 10 \log \frac{I}{I_0}$$

$$I = I_0 e^{-\mu x}$$

$$\mu_m = \frac{\mu}{\rho}$$

Electronics

Resistors

Preferred values for resistors (E24)
Series: 1.0 1.1 1.2 1.3 1.5 1.6 1.8 2.0 2.2
2.4 2.7 3.0 3.3 3.6 3.9 4.3 4.7 5.1 5.6 6.2
6.8 7.5 8.2 9.1 ohms
and multiples that are ten times greater

$$Z = \frac{V_{\text{rms}}}{I_{\text{rms}}}$$

$$\frac{1}{C_T} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$$

$$C_T = C_1 + C_2 + C_3 + \dots$$

$$X_C = \frac{1}{2\pi fC}$$

Alternating Currents

$$f = \frac{1}{T}$$

Operational amplifier

$$G = \frac{V_{\text{out}}}{V_{\text{in}}} \quad \text{voltage gain}$$

$$G = -\frac{R_f}{R_1} \quad \text{inverting}$$

$$G = 1 + \frac{R_f}{R_1} \quad \text{non-inverting}$$

$$V_{\text{out}} = -R_f \left(\frac{V_1}{R_1} + \frac{V_2}{R_2} + \frac{V_3}{R_3} \right) \quad \text{summing}$$

Turn over for the first question

Turn over ▶

SECTION A: NUCLEAR INSTABILITY

Answer **all** of this question.

- 1 (a) X and Y are two different β emitting sources. Initially they contain the same number of unstable nuclei. Both sources have their emissions recorded over a period of time. The *decay constant* of source X is greater than that of Y. State what is meant by decay constant and describe **two** differences in the recordings from the two sources.

You may be awarded additional marks to those shown in brackets for the quality of written communication in your answer.

.....

.....


.....

.....

.....

(3 marks)

- (b) The activity of a sample of radioactive iodine, $^{131}_{53}\text{I}$, is presented in the following graph.



- (i) Show that the decay constant of $^{131}_{53}\text{I}$ is about $1 \times 10^{-6} \text{ s}^{-1}$.

.....

.....

.....

.....

.....

- (ii) Calculate the half-life of $^{131}_{53}\text{I}$ in days.

.....

.....

.....

- (iii) Calculate the initial number of $^{131}_{53}\text{I}$ atoms in the sample.

.....

.....

.....

.....

(7 marks)

Turn over for the next question


| |
|-----------|
| |
| 10 |

SECTION B: MEDICAL PHYSICS

Answer **all** questions.

- 2 (a) **Figure 1** shows the ECG waveform produced when electrodes are attached to the chest of a healthy person. Label the axes with suitable scales and units.

Figure 1



(2 marks)

- (b) State what is meant by *depolarisation* and *repolarisation* and, in terms of ion movement, explain how each effect is caused.

depolarisation:

.....

.....

repolarisation:

.....

.....

(3 marks)

(c) State how the action of the atria and ventricles correspond to the waveform PQRST shown in **Figure 1**.

.....

.....

.....

.....

.....

.....

.....

.....

(3 marks)

| |
|----------|
| |
| 8 |

Turn over for the next question

Turn over ▶

3 Modern diagnostic X-ray tubes are used to produce sharp images of parts of the human skeleton.

(a) Explain how the design of the anode allows a sharp image to be produced.

.....
.....
.....
.....

(2 marks)

(b) State and explain **three** methods used to minimise the patient dose during diagnostic radiography.

1

.....

.....

2

.....

.....

3

.....

.....

(3 marks)

- (c) A beam of 50 keV X-ray photons is incident on and travels through a piece of bone of thickness 1.2 cm. Calculate the ratio of the transmitted intensity to incident intensity of the X-ray beam.

half-value thickness of bone for 50 keV X-ray photons = 1.5 cm

.....

.....

.....

.....

.....

.....

.....

.....

.....

(3 marks)

| |
|---|
| |
| 8 |

Turn over for the next question

Turn over ▶

- 4 (a) For each of the defects of vision listed below, describe the sharpness of the image formed of an object placed at a distance of 25 cm from the eye.

Slight hypermetropia

.....
.....

Slight astigmatism

.....
.....

(2 marks)

- (b) A converging lens is used to view an object placed 36 cm from it. A virtual image is formed which is 2.5 times larger than the object.

(i) State the defect of vision that this lens would be used to correct.

.....

(ii) Calculate the power of the lens.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

(4 marks)

| |
|---|
| 6 |
|---|

- 5 (a) Describe how the vibrations of a sound wave are transmitted from the outer ear to the inner ear. You should include an explanation of how the pressure changes are amplified.

You may be awarded additional marks to those shown in brackets for the quality of written communication in your answer.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(4 marks)

Question 5 continues on the next page

- (b) The dBA scale is used as a measure of perceived loudness which matches the response of the human ear. State the **two** main features of the dBA scale which enable it to match the response of the ear.

.....

.....

.....

.....

(2 marks)

| |
|---|
| |
| 6 |

Quality of Written Communication *(2 marks)*

| |
|---|
| |
| 2 |

END OF QUESTIONS

There are no questions printed on this page

There are no questions printed on this page